

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

1893



U. S. DEPARTMENT OF AGRICULTURE

FARMERS' BULLETIN No. 1732 *rev.*

Jan. 1940

GROWING BARLEY *for* MALT AND FEED



BARLEY IS GROWN for both malt and feed. It is the purpose of this bulletin to make available to the farmer such information as may enable him to decide which variety to grow and whether or not to exercise the care necessary to obtain grain that meets the maltster's requirements.

The maltsters of the Central and Eastern States desire such barleys as Oderbrucker, Manchuria, Wisconsin Barbless (Pedigree 38), Velvet, etc. They wish these barleys to be well grown, plump, mellow, carefully threshed, and free from disease and weather damage. Certain sections discussed in the bulletin are particularly adapted to the growing of these varieties and to the production of the quality desired. In the Western States varieties with larger kernels, such as Atlas, are grown. Barley of good malting quality is rarely produced except in sections where it is the major crop.

The greater part of the barley produced is certain to be fed. Barley is a very useful grain feed, and the yield per acre is high in the Northern and Western States. Many farmers should grow barley for consumption on the farm.

GROWING BARLEY FOR MALT AND FEED

By H. V. HARLAN, *principal agronomist*, and G. A. WIEBE, *agronomist*, *Division of Cereal Crops and Diseases, Bureau of Plant Industry*

CONTENTS

	Page		Page
Recent trends and present problems in barley growing.....	1	Varieties of barley—Continued.....	
Malting barley.....	2	Two-rowed.....	10
Northern and Eastern States.....	2	Smooth-awned.....	10
Central Plains.....	6	Trebi.....	10
Northern Plains.....	7	Coast and Atlas.....	11
Mountain States.....	7	Club Marliout.....	11
Pacific coast.....	8	Winter barleys.....	11
Southern States.....	8	Where to procure seed.....	11
Feed barley.....	8	Diseases.....	11
Varieties of barley.....	10	Commercial fertilizers.....	13
Manchuria, Oderbrucker, Odessa, and O. A. C. 21.....	10	What, when, and how much to seed.....	13

RECENT TRENDS AND PRESENT PROBLEMS IN BARLEY GROWING

BARLEY GROWS WELL in a cool, humid climate. It will stand more heat under semiarid conditions than under humid conditions. When grown in warmer climates, it is often winter-sown. The most favorable growing conditions are found in the Northern and Western States, where it produces a high yield of grain per acre. Gradually farmers in these States have come to realize the high value of the barley crop, and there has been a slow expansion of the acreage. The average yearly production now is much greater than before the prohibition era. Any increase between 1918 and 1933 was obviously for feed. If the present trend is continued, less than one-fourth of the crop will be absorbed by the maltsters. Farmers are naturally attracted by the premium for malting barley. It is a new source of revenue and a profitable one. Three-fourths of the average crop, however, will not receive a premium. In favorable years barley of good malting quality is produced over wide areas, and the price is accordingly depressed. The gains in balanced husbandry should not be sacrificed in mass attempts to produce malting barley. Only those farmers who have a reasonable chance of harvesting grain of the desired quality are justified in changing from a feed to a cash crop. The farmers most likely to profit from the new market are those located on suitable lands in barley-growing sections. In such locations varieties wanted by the buyers can be grown with little or no sacrifice of yield. The local market also is built around the malting demand, and the local grain dealer is in touch with the market and equipped to serve the grower by not allowing the better barley to lose its character through mixture in the elevator or in shipment. Facilities for preventing mixture are not available in localities where little barley is grown.

In general, feed barleys should be grown by farmers who expect to use the crop primarily as feed; by farmers in sections where little barley is grown; by farmers in sections where the malting varieties yield decidedly less than the feed varieties; and by farmers in any section where farms are of such character that good malting quality is rarely produced. The market preference for varieties is of little importance to the farmer who feeds his crop, and malting premiums are of little interest to the farmer who cannot grow malting barley.

Barley growers are at the crossroads. The growing of barley should be continued on a healthy basis free from violent shifts of acreage on account of large premiums or lack of premiums for malting barley. This can be done only by maintaining a large acreage destined for feed. The soundness and permanence of the barley acreage is dependent on the combined judgment of the individual planters.

MALTING BARLEY

NORTHERN AND EASTERN STATES

Any discussion of malting barley is complicated by the great diversity of soil and climate found in the United States. The types




FIGURE 1.—The three general regions of barley production in the United States. The heavy broken lines divide the country into the humid-spring region (northeast), the humid-winter region (southeast), and the semiarid region (west). The dots show the acreage according to the census of 1930. Localities suitable for barley growing are indicated by the present distribution of the crop.

suited to the several regions are so different and the methods of growing and caring for the crop so various that the subject cannot be discussed from the standpoint of a single section. The country, however, can be divided into regions which will be discussed separately. The three most important divisions are shown in figure 1. As far as the American market is concerned, the most important region consists of the Northern States east of the Missouri River.

Although many kinds of barley are grown in this region, the so-called "malting varieties" do well throughout the area, and an acceptable quality results where and when the soil and seasonal conditions are favorable. The malting market, however, is a critical one, and if the farmer wishes to be successful in this field he must be familiar with the requirements.

WHAT THE MALTSTERS WANT

The eastern maltsters and brewers want a plump, mellow, small-kerneled barley with firm hulls not easily damaged. They wish this barley to be all of the same type, sound, carefully threshed, and free from weather and disease damage. In actual field culture grain is seldom produced that is free of all objectionable features. The farmer growing barley for market should not only make every reasonable effort to enhance the quality as far as it is determined by the factors under his control, but should also be sufficiently familiar with malting requirements to be able to decide whether or not his barley meets these requirements.

Federal grades for malting barley under the Grain Standards Act have been established for barley grown east of the Rocky Mountains, effective July 1, 1937. The requirements for malting barley include the following specifications:

"This subclass shall include six-rowed barley of the class barley (class I) which meets the requirements of grades Nos. 1 to 3, inclusive, which, after the removal of dockage, contains not more than 5 percent of two-rowed and/or other types or varieties of barley of unsuitable malting type, such as Trebi and black; which contains not more than 15 percent of barley and other matter that will pass through a 20-gage metal sieve with slotted perforations 0.076 ($47/8_{64}$) of an inch wide and three-fourths of an inch long; which contains not more than 5 percent of skinned and/or broken kernels; which contains not more than 4 percent of damaged barley; and shall not include bleached barley. Barley of this subclass shall contain 75 percent or more of mellow barley kernels, which kernels are not, en masse, semisteely."

Black barleys (such as Lion) are objectionable not because they differ much from Oderbrucker, but because black kernels are sometimes difficult to distinguish from heat-damaged kernels and because the maltsters do not want them and pay less for barley when they are present.

Broken kernels are objectionable in barley intended for malting purposes because the broken kernels will not malt and must be removed to be sold as feed barley. Skinned kernels are even more objectionable than broken kernels because they cannot be screened out of barley intended for malting purposes and because they do not germinate and convert properly during the malting process. Frayed ends are objectionable. In many cases such kernels are damaged. Even if no real injury exists, malt made from barley containing many frayed ends has a poor appearance and will not sell so well as malt made from barley with unbroken hulls. Only 4 percent of damaged barley of all kinds is allowed in the malting grades. Damaged barley includes blighted, moldy, heat-damaged, sprouted, and badly weather-

damaged kernels. Blighted kernels, which include those affected by scab and stripe, may be very objectionable, and the limit of tolerance cannot be much greater than the amount specified in the grades.

LOCATION AND SOIL

Most of the barley that is malted is produced in a few States in the upper Mississippi Valley (fig. 2). In these States the best of the crop comes from a few localities. In the most-favored localities grain of malting quality is produced almost every year. Other sections produce good grain in 3 years out of 4, in half the years, or only occasionally. Small areas where high-quality barley is produced are to be found in all States from western New York to South Dakota. Such areas are well known to local farmers. Obviously the growers so favorably situated now produce and probably always will produce the greater part of the market requirements.



FIGURE 2.—A field of spring barley in the upper Mississippi Valley.

In the barley-growing areas some farms are better suited than others for the growing of high-quality barley. The best barley soils are well-drained loams. Barley cannot stand "wet feet." For this reason, heavy, poorly drained soils do not produce good barley in regions of frequent rains. Not only is the grain likely to be coarse, but the plants are more likely to be infected with blight and scab. On the other hand, light sandy soils are poor barley soils. Growth is not maintained at a sufficiently uniform rate, and ripening is often hastened by drought.

VARIETIES

The choice of variety is important. No variety, however, will produce malting barley in a section not suited to barley, nor can it overcome an adverse season. The fact that a variety produces grain of fine quality in western New York should not encourage a farmer in Iowa to plant it in his field. The State experiment stations are testing all promising sorts and are the best sources of information for local use. Full information can be had on many varieties, but some of the newer ones must be grown several years before they can be accurately appraised. The type wanted is definite. Maltsters ex-

press a decided preference for barleys of the Manchuria-Oderbrucker group and smooth-awned hybrids from these parents. Large-kernelled six-rowed varieties and two-rowed sorts are apparently not in demand in this section. Oderbrucker, Manchuria, Odessa, and the smooth-awned hybrids, Wisconsin Barbless (Pedigree 38), and Velvet, seem to be acceptable. Manchuria, which is blue or mixed blue and white, and such blue selections of Manchuria as O. A. C. 21, are disliked by some maltsters because of the blue color, which makes the determination of mellowness more difficult. The blue color in itself is entirely harmless, being merely a pigment deposited in the outer layer of the endosperm. It has nothing to do with malting quality. The smooth-awned varieties have a tendency to peel more easily than Manchuria or Oderbrucker. This is an undesirable character. It is worse in poorly grown barley than in plump, well-matured barley. When carefully threshed, the smooth-awned sorts meet the market requirements.

SEED AND SEEDING

Seed should be sound, clean, and of high germination. The use of the fanning mill is important, because it removes weed seeds and many of the diseased kernels. There is a common impression that seed of high malting quality is desirable for sowing. This is true only as far as malting quality indicates strong germination and freedom from disease. Hard, flinty, steely barley is just as desirable for seeding as is the finest malting barley of the same variety if the germination is equal.

Seeding should be early for the locality. Barley does not grow well in hot, humid weather, and there is a decided advantage in early seeding.

ROTATION

Barley of high quality is seldom produced on land that is too highly fertile. On rich lands manures are best applied to the preceding crop. The residual effect is sufficient for barley. Coarse grain and lodging are often prevented in this way. In areas where scab is serious, barley should preferably follow some other crop than corn or small grain.

HARVESTING AND STACKING

In the harvesting of malting barley the farmer must meet a number of serious problems. Sound, fully ripened grain is without question of great importance to the maltster. On the other hand, the longer the grain stays in the field, the greater the hazards from storm and moisture damage. Ripened under the best of conditions in a humid climate, the grain assumes a bright straw color.

Quality is often seriously impaired during ripening and harvesting. The crop should stand until it is fully ripe. Not only is the grain more mellow, but the proteins cause less trouble in brewing. The proteins of growing grain are very different from those in the fully matured kernel. The same is true, in a way, with the starch. If growth is arrested by drought or too early cutting, the kernels are likely to be "steely." Such grain does not malt well and brews worse. Care should be taken not to cut the crop too soon. Probably

the thumbnail test to determine maturity is as good as any. Grain is ready to cut when it has reached the point where it will retain the imprint of the thumbnail pressed against it.

Color can easily be overemphasized. In dry sections or where the crop ripens without rain or dew, perfectly matured grain may have very little of the yellow color that is characteristic of grain ripened where light rains and dews are common. A deeper color in humid sections is not a cause for a lower price, and the farmer should not sacrifice quality to secure grain that exhibits the same brightness as that found in the drier sections.

Frequent rains at harvest time often injure the quality of the grain. Any considerable number of kernels with blackened ends lowers the market value. Musty odor is a danger signal to the barley buyer, and even a small percentage of decayed kernels may remove a shipment from the malting class. Extra care must be used in shocking and stacking barley intended for the malting market. Minimum exposure to unfavorable weather is desirable.

Stacking improves the quality of barley. Something occurs during the process of going through the sweat in the stack that makes barley more mellow and increases both the vigor and percentage of germination.

THRESHING

Great care should be exercised in threshing malting barley. Skinned and broken kernels are a hazard to the maltster, and any chance of receiving a premium for the crop can be ruined in threshing. The maltster would prefer to have a few awn fragments remaining on the grain than to be confronted with threshing damage. Especial care may need to be taken with the smooth-awned varieties. Any thresher can adjust his machine to decrease threshing damage. Square teeth in the concaves instead of rounded ones result in fewer broken kernels in the final product. The elimination of end play in the cylinder is of great importance.

CENTRAL PLAINS

The area discussed as suitable for the Manchuria-Oderbrucker type of barley has been very crudely bounded. "East of the Missouri," of course, does not mean that the Missouri River marks a definite boundary. Eastern Nebraska, for instance, is as much a part of the region as southern Iowa. As one goes farther west, however, conditions gradually change. The Manchuria-Oderbrucker types do not yield so much as many others do, and they shatter badly in the drier sections.

The western types come into competition with the eastern ones in northwestern Kansas and adjacent areas. The most dependable varieties in this transition area in the past have been related to those long grown on the Pacific coast. Manchuria-Oderbrucker types also can be grown, although at some sacrifice of yield. The maltsters on the eastern markets prefer this latter type, but when of good quality the larger-kerneled Coast type is also in some demand. Under such conditions the farmer has a difficult decision to make. The correctness of this decision is dependent on many factors beyond his control. Whether he has good barley depends on the season. Whether or not

there will be a profitable premium for it depends on the seasonal conditions over a wide area.

NORTHERN PLAINS

Recommendations for the dry lands of Montana and adjacent areas are equally difficult at present. The Manchuria-Oderbrucker barleys do not yield well, and they shatter badly. The brewing value of Montana-grown Horn and Trebi has not been established. The quality exhibited by these varieties in other areas is little indication of what they may do here. When brewers have used barley produced in this area during 1 or 2 average years, a reasonably sound recommendation can be made. Until this information is available, none can be given.

MOUNTAIN STATES

The agriculture of the western mountain country, like that of all mountain countries, is largely confined to the many irrigated valleys, each one a problem in itself. Regardless of variety, the grain from these valleys will always be plumper, of lower nitrogen content, and more mellow than that produced in the upper Mississippi Valley. Grain of the character obtained in the upper Mississippi Valley could not be produced even if it were desired. Furthermore, in this area, barleys of the Manchuria-Oderbrucker type shed most of their kernels before harvest. The brewing problem here, then, becomes quite different from that in Milwaukee or Chicago. If the brewer uses local material, he will probably find it good and far more uniform than that of the East, but he must of necessity develop his own methods of utilization.

Volume is, of course, limited since under highly specialized irrigation farming, the percentage of land in grain is not large, and much of the barley produced is needed for feed. The scarcity of feed grains in the Mountain States and the high price paid for them often makes the importation of malt economical.

Since the better barley in the Mountain States is produced under irrigation, the judicious use of water is important. Unfortunately, there are no rules by which the most favorable conditions can be obtained or maintained. No two fields or two seasons are alike. The object of irrigation is, of course, to keep the plants growing well, and the timing of the watering must depend on the judgment of the farmer. Wherever possible, the land should be so handled that there will be sufficient water in the soil to germinate the grain. Applications of water after seeding to start growth are usually disastrous. Too rank growth at heading time is undesirable. Late irrigations are also a problem. There is danger of lodging the grain. On the other hand, failure of water before full maturity cuts heavily into the yield and lowers the quality. Barley kernels continue to grow until they dry. Any kernel that will not retain the imprint of the thumbnail is still growing. Thinly seeded nurse crops often produce high yields because irrigations are continued to save the grass. If possible, the last irrigation should be late enough to furnish moisture until the plant has ripened naturally. When malting barley is grown under irrigation, the threshing should be watched carefully. The dry

kernels break easily, and most threshers in this area are inexperienced in adjusting their machines to meet these particular market requirements.

PACIFIC COAST

The conditions on the Pacific coast are peculiar to that area. Most of the market barley is raised in the interior valleys of California. The small-kerneled Manchuria-Oderbrucker barleys cannot be grown. England furnishes the market, and her maltsters prefer barleys of the Coast type. Considerable barley is malted locally, and in some years fairly large quantities of California barley are shipped east for malting purposes. Atlas is the dominant variety at present. Purity of type and freedom from threshing damage are important and are under the control of the farmer. Club Mariout is useful in late sowing and, when well grown, is used by some of the English maltsters. Its price on the English market is less than that of Atlas. The Coast type of barley was formerly produced in some quantity in eastern Oregon and Washington. There are also special sections where barleys of other types and of high quality can be grown, as in the Salinas Valley of California and the section west of the Cascades in Oregon and Washington. Hannchen is the dominant variety in western Oregon and in the Klamath Falls district, which lies on the border between Oregon and California. This variety is used for malting both on the Pacific coast and in the East.

SOUTHERN STATES

There are many localities where barley can be grown that cannot be separately discussed. In the Southeastern States, for instance, bearded winter barley is grown that is quite similar to the Manchuria-Oderbrucker type. Conditions at harvest are often unfavorable, and over most of the South the growing of malting barley is not recommended.

FEED BARLEY

Feed barley is barley used as feed. It may or may not be of the varieties or qualities desired by maltsters. It may be grown solely for feed, or it may be the result of an unsuccessful attempt to grow barley acceptable to the maltsters.

Market terminology has done much to retard the expansion of the barley acreage. Grain rejected by the maltsters is called "feed barley." It usually consists of inferior grain and is only second-rate feeding material. Barley grown for feed need not be inferior grain. Indeed, in the good barley sections it is usually of relatively high quality.

In many ways it is unfortunate that feed barleys are not a crop entirely distinct from malting barleys. The value of barley for feeding must be measured by its usefulness on the farm. The return in terms of pounds of pork or gallons of milk should be the farmer's gage. The appraisal of feed barley would be much less difficult if there were no other barley market.

In the years when brewing was prohibited there was a marked increase in the barley acreage. This increase was, of course, for feed.

Much of the recent shifting in barley acreage has been outside the better-known area of malting-barley production. There was a short-lived expansion into the Corn Belt. A heavy infection of scab discouraged any rapid or careless expansion in this direction. On the Great Plains there is an apparently permanent increase. This has been particularly heavy in northwestern Kansas. In western Oklahoma and adjacent sections there has been a fluctuating acreage of winter barley with an upward trend. There is also an active interest in winter barley in the Southeastern States, where the acreage for feed is increasing. Spring barley is best adapted to the Northern and Western States. Much of this area is outside the Corn Belt. A grain feed that may be substituted for corn is, of course, of great value. Barley is particularly suited to this purpose, both because of its yielding capacity and its feeding value.

On favorable soils, barley yields more pounds per acre than any other small grain. It fits in well with the systems of agriculture practiced in most of the areas where the crop is adapted. It can be fed to all classes of livestock and is constantly increasing in popularity with dairy farmers. It is almost the equal of corn as a feed. As in the case of corn, it is especially prized as a feed for hogs, because it produces firm pork. The value of clean and sound barley for feeding hogs depends largely upon its weight per bushel. Barley weighing 48 pounds per bushel has almost the same value as corn. If it weighs only 44 pounds per bushel, experiments indicate that it is worth about 10 percent less per bushel for hog feeding. This is due to its higher fiber content, greater bulkiness and, therefore, lower nutritive value. For other classes of livestock the higher fiber content is not such an important factor. In fattening cattle, scabby barley, which has a considerably lighter weight per bushel than sound barley, has practically the same value pound for pound as sound barley. Ordinarily, hogs will not eat scabby barley.

There is little difference in the feeding value of varieties. The farmer growing barley for feed ordinarily should use the highest-yielding variety. Other things being equal, the percentage of hull is higher on small-kerneled than on large-kerneled barleys. Feeding tests, however, have shown little difference in the value of the different kinds if they are plump and well grown.

The barley kernel is too hard to use satisfactorily as feed without some previous preparation. Grinding is the most common treatment in the Central and Eastern States. The grain should not be ground fine. Fine-ground barley makes a pasty mass when chewed by animals, and feeding it results in reduced consumption. When grinding barley, the machine should be so set as merely to crack the grain rather than pulverize it. In the West much of the barley is rolled. This is an ideal method of preparation. A jet of steam softens and moistens the kernel so that little loose material results. At the same time the entire kernel is flattened to a soft, easily crushed disk.

When barley is to be mixed with chicken feed the hulls are sometimes removed. This separation need not be complete. The removal of most of the hull increases the palatability for chickens and greatly improves the quality of the feed by reducing the amount of roughage.

VARIETIES OF BARLEY

More than 5,000 varieties of barley have been tested by the United States Department of Agriculture and State experiment stations. Only a few of them are under cultivation on farms. There is a decided advantage in growing a limited number of varieties. Mixtures are a great disadvantage to the maltster, and the fewer sorts grown the less the confusion in marketing. Some of the most important varieties are briefly discussed below.

MANCHURIA, ODERBRUCKER, ODESSA, AND O. A. C. 21

Manchuria, Oderbrucker, Odessa, and O. A. C. 21 are similar in kernel character and are the type preferred by the maltsters of the Mississippi Valley. Although all four are grown in the Northern States east of the Missouri River, the centers of production are distinct. Odessa is largely confined to South Dakota. Oderbrucker is grown most extensively in Wisconsin. Manchuria is more common in western Minnesota and North Dakota, although a selection, Featherston, is grown to a limited extent in New York. O. A. C. 21 is extensively grown in Canada. The acreage of all these varieties has been reduced in the United States because of farm preference for the smooth-awned sorts. The grain of Oderbrucker is the whitest of the lot. Manchuria consists of a mixture of blue and white strains, whereas O. A. C. 21 is a blue strain of Manchuria.

TWO-ROWED

Two-rowed barleys are not grown extensively. Most brewers prefer a smaller-kerneled barley of higher diastatic power. Spartan is grown on a small acreage in Michigan and nearby States. Alpha is grown in New York and Pennsylvania. There are scattered fields of Hanna and Hannchen throughout the Plains States. Horn is common in Montana, and there is a remnant of Chevalier in a few localities, such as the Salinas Valley of California. Hannchen is grown in western Oregon and in the Klamath Falls district, which lies on the border between Oregon and California.

SMOOTH-AWNED

Farmers naturally prefer smooth-awned barleys. The Department of Agriculture and many State stations have tried to breed suitable varieties. The factor of smoothness is complex, and difficulty has been experienced in keeping such barleys uniformly smooth. Several sorts are sufficiently so for commercial purposes, and their yields have been satisfactory. Among those in common cultivation are Velvet, Glabron, and Wisconsin Barbless (Pedigree 38) in the upper Mississippi Valley. Comfort is grown in Nebraska. Vaughn and Hero have attained a small acreage in California. The former variety has produced high yields in Arizona and the southern plains. Flynn was recently released in western Kansas and is increasing in acreage.

TREBI

Sixteen years ago Trebi was released to farmers of southern Idaho. All that was known then was that it was well adapted to irrigated

lands in this section. Since that date it has spread over much of the northern part of the United States and up into Canada. Although its actual field cultivation has not extended east of the Mississippi River, it has produced high yields on experiment station plots as far east as the eastern part of Canada. The variety is unquestionably one of the most vigorous that has been tested. Despite its low resistance to disease, it has outyielded other varieties in sections to which by any other measure it seems not to be suited. The grain as produced in eastern North Dakota is very different from that of Idaho, but from the standpoint of feed, it deserves consideration. Maltsters of the Mississippi Valley do not like Trebi. Their acquaintance with the variety is limited to that grown in the northern plains and in the western half of the upper Mississippi Valley. Brewers of the Rocky Mountain area may find the Trebi grown in this area satisfactory. This is especially true of that produced under irrigation.

COAST AND ATLAS

Coast was the common barley of the Western States for many years. The original Coast and Atlas, a selection of Coast, are still grown on most of the acreage in California. Coast is common throughout the Western States.

CLUB MARIOUT

Club Mariout is limited mostly to California, where it is popular for late seeding.

WINTER BARLEYS

Tennessee Winter (fig. 3) and Wisconsin Winter are two of the hardiest winter barleys (fig. 4). They are also high-yielding sorts. Both are bearded. In the Southeastern States hooded varieties are more popular than bearded varieties. The first distribution of hooded winter barleys came from the Tennessee Agricultural Experiment Station. The varieties released from this station have been added to by other southern stations and by natural hybridization occurring on farms. The hooded varieties yield less than the old bearded Tennessee Winter but are preferred by farmers because of the greater comfort in handling. Missouri Early Beardless, a hooded sort, is grown extensively in Missouri and neighboring States. It is used for fall and spring pasture and for grain.

WHERE TO PROCURE SEED

Most of the varieties recommended are already in wide cultivation, and may be obtained from nearby farmers. Most States have seed-growers' associations from which certified seed can be obtained. The State experiment stations can usually furnish information as to sources. Many local seedsmen carry the more important varieties.

DISEASES

Four of the most important diseases of barley for which chemical or cultural treatments are known are scab, stripe, covered smut, and one kind of loose smut.



FIGURE 3.—A field of Tennessee Winter barley.




FIGURE 4.—A field showing winter barley as a nurse crop for red clover.

Scab is largely confined to the Corn Belt. This disease affects the quality of the grain and reduces the yield. Barley with more than a small percentage of scab cannot be used for malting. It is also unsuitable for feeding pigs, causing vomiting in extreme cases. If the grain is less heavily infected, the pigs fail to make normal gains. Scabby barley can be fed to all classes of cattle and sheep.

Treatments for scab are largely preventive. Since the disease is carried over on cornstalks and grain stubble, such material should be carefully plowed under before seeding. The disk obviously is no substitute for the plow on stubble land. Fields that were not planted to either corn or small grain the previous year are likely to have less scab. In scab areas the seed should be run through the fanning mill before being sown, to remove as many infected kernels as possible. Also the grain should be treated with a mercury-dust compound (New Improved Ceresan) for control of the seedling blight due to scab-infected seed. Follow carefully the directions for treatment given on the label of the can. *Mercury-dust compounds are poisonous. Treat seed outdoors or in a well-ventilated building. Avoid inhaling large quantities of the dust. Wear a dry cloth or dust mask over the nose and mouth. Do not feed treated seed to farm animals.*

Blighted barley is a market problem. Darkened, underdeveloped kernels are caused by a wide variety of organisms in addition to that of scab. The effect of many of these are less objectionable than scab, especially for hog feeding. The buyers are often unable to determine the cause of the damage or to distinguish scabby kernels from kernels affected by other organisms. Seed treatments and clean culture reduce the amount of infection of some of these organisms.

Covered smut is the most widely distributed of the important diseases of barley. The total loss due to it is considerable. The percentage of infection varies enormously with season and region. Where covered smut is common, clean seed should be used where possible. If clean seed is not at hand, the farmers should resort to dust treatment. Mercury dust (New Improved Ceresan) is a logical choice not only because of its usefulness in controlling covered smut but also because it controls stripe, seedling blight due to scab, and one kind of loose smut where these are present. The other kind of loose smut can be controlled by the hot-water treatment. Because of the serious seed injury that may result, the hot-water treatment is not recommended for use by farmers. County agricultural agents can give suggestions as to sources of chemicals and methods of treatment.

COMMERCIAL FERTILIZERS

Most of the barley crop is grown in sections where commercial fertilizers are not widely used. Individual farmers had best secure information on fertilizers from their State experiment stations and substations.

WHAT, WHEN, AND HOW MUCH TO SEED

The recommendations that follow as to varieties to seed, time of seeding, and quantity to seed are in accord with those of the State experiment stations as far as information is available.

Arizona.—For the irrigated valleys of southern Arizona, Arivat Vaughn, Common Six-Row, and Scarab are the recommended varieties. For higher elevations of 3,500 feet, or more, Trebi should be used. Where barley is intended for pasture, it should be seeded early in the fall, when danger from hot weather is past. This is about early October for the Salt River Valley. For grain purposes seeding should be done from late November to early January. At high elevations (6,000 feet) spring barley should be seeded April 15 to May 15. The seeding rate for pasture is 90 pounds per acre; for grain, 75 pounds per acre.

Arkansas.—Selection 6, a barley of the Union Winter type, and the hooded varieties, Tennessee Beardless and Missouri Early Beardless, are suggested for general use in Arkansas. It is recommended that barley be sown the first week in October at the rate of 7 pecks per acre.

California.—Atlas is recommended for the production of malting barley. Vaughn is the highest yielding variety and should be grown for feed. Where seeding cannot be done until late, Club Mariout is suggested. The best time to seed is from late October until mid-January. A rate of 70 to 90 pounds per acre is sufficient, when a drill is used. When the seed is to be broadcast, however, the rate should be increased 25 percent.

Colorado.—For irrigation conditions similar to those at Fort Collins the recommended varieties are Lico, Trebi, Colsess, and Hannehen. If the crop is intended for malting, Wisconsin Barbless (Pedigree 38) and Velvet are recommended. For high-altitude conditions similar to those at Fort Lewis, Trebi and Colsess are recommended. Colsess is particularly well-suited as a nurse crop for alfalfa and red clover and for hay production. For dry-land conditions similar to those at Akron, the recommended varieties are Club Mariout, Flynn, and Vance Smyrna. In areas under irrigation, seeding should be done April 1 to 20 and progressively later at higher elevations. A good seeding rate is 2 bushels an acre. For dry land, seedings should be done in late March or early April at a rate of 4 pecks an acre.

Georgia.—The most promising variety appears to be Greece. At Experiment and Tifton, Texas Winter and Tennessee Winter Selection 66 produce high yields. Barley should be seeded by early October in the northwestern part of the State and by October 25 elsewhere. The recommended seeding rate is 6 to 8 pecks per acre.

Idaho.—For the irrigated lands and the Palouse section Trebi is the recommended variety. In the cut-over lands of northern Idaho, Hannehen and Union Beardless are the recommended varieties. For fall seeding in the Palouse section, Winter Club should be used. In the irrigated sections seeding should be done in late March or early April at the rate of 100 pounds an acre. In the nonirrigated sections the time of seeding is about the same, and the rate of seeding is 6 pecks an acre, decreasing, in the drier areas, to 4 pecks.

Illinois.—Wisconsin Barbless (Pedigree 38) is the recommended variety for general production for both market and feed. Where Trebi is grown, it should be considered only as a feed barley. Spartan is useful where a stiff-strawed variety is needed. On the southern edge of the barley section early seeding is very important. Around Urbana barley should be seeded in late February or early March. In the northern part of the State it may be seeded somewhat later. The recommended seeding rate is 8 pecks an acre. Some winter barley is grown in southern Illinois. Purdue 21, Purdue 1101, Kentucky No. 1, and Missouri Early Beardless are the best adapted varieties. Winter barley should be seeded in September at the rate of 6 to 8 pecks an acre.

Indiana.—Only limited sections in Indiana are suited to the cultivation of barley. Wisconsin Barbless (Pedigree 38) and Velvet are recommended. Spring barley should be seeded as early as possible at the rate of 2 to 2½ bushels an acre. September is the most favorable time for fall seeding.

Iowa.—Velvet and Wisconsin Barbless (Pedigree 38) are the recommended varieties for malting or feed. Barley responds favorably to early seeding. It should be sown no later than the middle of April. Seedings made after this date yield much less. A satisfactory seeding rate is 2 bushels an acre.

Kansas.—For spring sowing Flynn is the recommended variety. Barley should be sown between March 15 and April 10. The best rate of seeding is from 5 to 7 pecks an acre, depending on the locality. Where winter barley is sown, locally adapted strains should be used. Fall sowing should be done between September 15 and 30, at the rate of 2 bushels an acre.

Kentucky.—Winter barley is a fairly dependable crop on the better soils in Kentucky. Local strains, which appear to be identical with Tennessee Winter,

and Kentucky No. 1 are the most widely grown varieties. Kentucky No. 1 has a stiffer straw than the ordinary types. Barley should be sown October 1 in the northern half of the State and October 10 in the southern half. The rate of seeding varies from $1\frac{1}{2}$ to 2 bushels an acre, depending on soil productivity and time of seeding.

Maryland.—Winter barley predominates in Maryland. Tennessee Winter and Wisconsin Winter are good varieties for general use. Marnobarb is preferable except at higher altitudes. Barley should be sown by the last of September. The recommended seeding rate is 2 bushels an acre.

Michigan.—The present trend is away from two-rowed barleys toward the six-rowed type. Wisconsin Barbless (Pedigree 38), a six-rowed type, is the recommended variety for market and feed. Where barley is used principally for feed or as a nurse crop, Spartan is a good variety to use. It is particularly suited to this latter use because of its early maturity, stiff straw, and sparse foliage. Barley should be seeded as soon as the ground can be properly prepared in the spring, at the rate of $1\frac{1}{2}$ to 2 bushels an acre.

Minnesota.—The varieties recommended for all sections of the State are Wisconsin Barbless (Pedigree 38), Velvet, Glabron, Peatland, and Improved Manchuria, C. I. No. 2330. Peatland is recommended particularly for peat lands or in regions where scab is a serious problem. Where malting barley is grown, Wisconsin Barbless (Pedigree 38), Velvet, Improved Manchuria, and Peatland should be used. Glabron should be grown only for feed. Barley should be seeded as early as the ground can be prepared. This can usually be done by April 10 in some parts of the State, whereas in others it may not be possible before April 25. The recommended seeding rate is 2 bushels an acre.

Missouri.—Winter barley is used extensively for fall and spring pasture and for feed. It is an excellent nurse crop for grasses and legumes and for soil-erosion control. For these purposes the hooded variety, Missouri Early Beardless, is generally recommended. It has not produced as much grain as some of the bearded varieties, nor is it as winter hardy as some other varieties. It is, however, usually sufficiently hardy to survive the winters except at the northern edge of the State, where it frequently winter-kills. When intended for pasture, barley should be sown in late August or early September at a rate of 8 pecks an acre. For grain purposes the best results are obtained when it is sown in late September or early October at a rate of 6 to 8 pecks an acre. Spring varieties should be sown in March at the rate of 8 pecks an acre.

Montana.—Trebi and Horn are the recommended varieties. For dry land, seeding should be done as early as the season will permit and preferably on clean summer fallow at the rate of 4 to 5 pecks an acre. For irrigated lands, 8 pecks an acre is a good seeding rate, and the best time to seed is May 1 to May 15.

Nebraska.—Spartan, Trebi, Short Comfort, and Club Mariout are the recommended varieties. Trebi is especially recommended for the irrigated lands in the North Platte Valley. Barley is usually sown at the rate of 2 to $2\frac{1}{2}$ bushels an acre in eastern Nebraska, the rate diminishing materially toward the west under nonirrigation conditions. It is recommended that barley be sown in late March or early April in the eastern part of the State and progressively somewhat later toward the west.

Nevada.—Trebi is the recommended variety for most of Nevada, except on lighter soils, where Coast should be used. April is the most favorable time to seed in the north and central part of the State. In the southern part seeding should be done somewhat earlier. Barley is usually sown 2 to 4 weeks after spring wheat. The recommended rate is 80 to 90 pounds an acre.

New Jersey.—Velvet and Comfort are recommended for spring seeding, and Marnobarb where a fall-sown variety is desired. Spring-sown varieties should be sown in mid-April at the rate of 8 to 10 pecks an acre. Fall-sown barley should be sown between September 20 and October 1. The recommended rate is 6 to 8 pecks an acre.

New Mexico.—Trebi and Club Mariout are the varieties recommended to grow under irrigation in the Rio Grande Valley near El Paso and Albuquerque and in the Pecos Valley from Roswell to Carlsbad. Barley should be seeded February 1 to 15 at the rate of 90 to 100 pounds an acre. For dry-land conditions similar to those at Capulin, White Smyrna and Stavropol should be used. Seeding should be done late in May or early in June at the rate of 1 bushel an acre.

New York.—The recommended varieties for New York are Alpha for feed, either grown alone or mixed with oats, and Wisconsin Barbless (Pedigree 38)

as a market barley. Barley should be sown in April at the rate of 2 bushels an acre.

North Carolina.—Tennessee Beardless No. 6, a hooded sort, is the recommended variety for North Carolina. Barley should be sown during October. A satisfactory seeding rate is 2 bushels an acre.

North Dakota.—Trebi is recommended as a feed barley for most of North Dakota. On farms where market barley can be grown Manchuria, Velvet, and Wisconsin Barbless (Pedigree 38) are recommended. Odessa, a variety popular in South Dakota, shows considerable promise. Steigum and Horn show promise in the drier sections in the western part of the State. In the eastern part, Manchuria, Velvet, and Wisconsin Barbless (Pedigree 38) should be seeded at the rate of 6 pecks an acre. Farther west 5 or 6 pecks is sufficient. For Trebi and other large-seeded types, the rate should be somewhat higher. Barley should be seeded in late April or early May.

Ohio.—Spring barley is grown only in northwestern Ohio. Here Velvet and Glabron are the recommended varieties and should be sown as early as possible. Winter barley is grown in the central and southern part of the State. Hardy strains of Tennessee Winter are recommended. Winter barley should be seeded from the middle to the last of September. The rate of 2 bushels an acre is recommended for both spring and fall seeding.

Oklahoma.—Where barley is fall-sown, Ward is the recommended variety for the Panhandle and the northern tier of counties, whereas in other parts of northwestern Oklahoma, Wisconsin Winter and Tennessee Winter should be used. For conditions similar to those at Lawton, the varieties Michigan, Wisconsin Winter, or locally adapted bearded sorts are recommended. Barley should be seeded by October 15. For pasture early seeding is desirable. A good seeding rate is 2 bushels an acre. Atlas and White Smyrna are the best where spring barley is grown and should be sown February 10 to March 1. The rate varies from 5 pecks in the western section to 8 pecks on the better lands, where the rainfall is heavier.

Oregon.—For western Oregon Hannchen and Union Beardless are recommended for spring seeding. The latter variety is especially suited for fertile soil because of its stiff straw. Where winter barley is grown, Santiam, O. A. C. No. 1, and O. A. C. No. 6 are recommended. In this section fall-sown barley should be seeded October 10 to 30; spring-sown barley, March 15 to April 15. The rate varies from $1\frac{1}{2}$ to $2\frac{1}{4}$ bushels an acre.

For conditions similar to those near Pendleton, Trebi is recommended. Meloy should be used where a hooded variety is desired for hay. Barley should be seeded March 1 to 15 at the rate of 2 bushels an acre.

Under the limited-rainfall conditions in the Moro section, Club Mariout is the most widely grown variety, although Flynn and Meloy are also popular, especially when cut for hay. In this section barley should be seeded as early in the spring as the ground can be prepared, at the rate of $1\frac{1}{4}$ to $1\frac{3}{4}$ bushels an acre.

At Union, Trebi should be used for grain, and Union Beardless should be grown where barley is cut for hay or on rich ground where stiffness of straw is a desirable factor. Seeding should be done March 15 to April 15 at the rate of $1\frac{1}{2}$ to 2 bushels an acre.

In the Burns section, Trebi, O. A. C. No. 7, and Union Beardless are the best varieties to grow. Barley should be seeded in this section about the first of May at the rate of 2 to $2\frac{1}{4}$ bushels an acre.

Pennsylvania.—For spring seeding Wisconsin Barbless (Pedigree 38) and Alpha are recommended. Seeding should be done about April 10 in the southeastern part, about April 24 in the State College section, and about May 8 in the northern counties. The recommended seeding rate is 2 bushels an acre. For fall seeding Kentucky No. 1, Tennessee Winter, and a local strain from Cumberland County are the most promising. Winter barley is an uncertain crop north of the center of the State. Fall-sown barley should be seeded the first week of September in the State College section and 2 to 3 weeks later in the southeastern counties. The usual seeding rate is 2 bushels an acre.

South Carolina.—Hooded or awnless varieties are favored by farmers in South Carolina. The recommended varieties are Beardless Winter and Bearded Winter. Barley is often used for fall and winter pasture or as a cover crop. October is the most favorable month for seeding, and $1\frac{1}{2}$ to 2 bushels an acre should be sown.

South Dakota.—Where barley is intended for malting, Odessa, Velvet, and Wisconsin Barbless (Pedigree 38) are the recommended varieties. Wisconsin Barbless should be sown only in the southeastern section. Where the grain is intended for feed the recommended varieties are Trebi, Glabron, and Spartan. In the drier sections, Horn, White Smyrna, and Ace are satisfactory varieties to grow. In the irrigated sections in the western part of the State, Trebi should be grown. Barley should be seeded early. Sowings made later than April 15 are likely to result in reduced yields. It is recommended that 6 pecks an acre be sown in the eastern part and 5 pecks in the central and western parts of the State. In irrigated sections 8 pecks should be sown.

Tennessee.—The recommended varieties for Tennessee are Tennessee Winter Selection 52, Union Winter, and Tennessee Beardless 5. The latter variety is a hooded sort and is preferred by some farmers. Barley should be seeded the latter part of September or in early October. A seeding rate of 8 pecks an acre is recommended. Under favorable conditions slightly less seed of Tennessee Winter Selection 52 and Union Winter might be used but not of Tennessee Beardless 5.

Texas.—Wintex, Finley, Texas Winter, and other intermediate winter strains of Tennessee Winter are recommended for fall planting (October 1 to 20) in central and northeastern Texas. The intermediate winter types also are more satisfactory for very early spring planting (January 1 to 20) in this section and westward than the true spring varieties like Vaughn and Stavropol. In extreme southwest Texas the true spring varieties are recommended for fall planting, as they rarely winter-kill and they mature ahead of hot weather and rust. A seeding rate of 2 bushels an acre is recommended for central and northeastern Texas, whereas 6 pecks or slightly less should be used in west Texas and the Panhandle.

Utah.—Velvon and Trebi are the recommended varieties for the irrigated lands of Utah. They should be seeded in early spring at the rate of 2 bushels an acre. Where a winter variety is desired, Bulgarian and Winter Club are suggested for the Nephi and Logan sections, respectively. Winter barley should be seeded in September or October at the rate of 7 pecks an acre.

Virginia.—The varieties generally recommended for Virginia are Wisconsin Winter and Tennessee Winter. Barley should be seeded September 15 to 30 at the rate of 2 bushels an acre.

Washington.—Beldi Giant, Rufflyn, and Blue are satisfactory six-rowed varieties for conditions similar to those at Pullman. Haunchen is the recommended two-rowed variety and Horsford where a hooded strain is desired. Winter Club is recommended as a winter variety. Under conditions of limited rainfall similar to those near Lind, Beldi Giant and Rufflyn are the best varieties to sow. Barley should be seeded as early in the spring as conditions will permit. In the drier sections, seeding should be done in March or early April. In the better-watered part of the Palouse April is the most satisfactory time. Winter barley should be seeded early in September. Seeding rates vary from 5 pecks an acre in the drier soils to 8 pecks in the best of the Palouse.

West Virginia.—Alpha is the most promising spring variety. It should be seeded in April at the rate of 6 pecks an acre. Limited tests with fall-sown varieties show that Kentucky No. 1 and Scottish Pearl are the most dependable. Fall-sown varieties should be seeded August 25 to September 5 at the rate of 8 pecks an acre.

Wisconsin.—Wisconsin Barbless (Pedigree 38), Oderbrucker, and Velvet are the recommended varieties for Wisconsin. Barley is usually sown after corn or oats. The seeding time varies from April 1 to May 7, but the average is about the third week in April. The recommended rate is $1\frac{1}{2}$ bushels an acre, except for Oderbrucker, 2 bushels of which should be used.

Wyoming.—Coast and Trebi are recommended for dry land. Where the straw is to be utilized, Horn, Smyrna, and Vaughn are suggested. For irrigated land, Horn and Trebi are satisfactory. Five pecks an acre is the recommended seeding rate on dry land and 8 pecks on irrigated land. The best time to seed is from April 15 to May 1.

**ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE
WHEN THIS PUBLICATION WAS LAST PRINTED**

<i>Secretary of Agriculture</i>	HENRY A. WALLACE.
<i>Under Secretary</i>	M. L. WILSON.
<i>Assistant Secretary</i>	HARRY L. BROWN.
<i>Director of Information</i>	M. S. EISENHOWER.
<i>Director of Extension Work</i>	C. W. WARBURTON.
<i>Director of Finance</i>	W. A. JUMP.
<i>Director of Personnel</i>	ROY F. HENDRICKSON.
<i>Director of Research</i>	JAMES T. JARDINE.
<i>Director of Marketing and Regulatory Work</i>	A. G. BLACK.
<i>Solicitor</i>	MASTIN G. WHITE.
<i>Land Use Coordinator</i>	M. S. EISENHOWER.
<i>Office of Plant and Operations</i>	ARTHUR B. THATCHER, <i>Chief</i> .
<i>Office of C. C. C. Activities</i>	FRED W. MORRELL, <i>Chief</i> .
<i>Office of Experiment Stations</i>	JAMES T. JARDINE, <i>Chief</i> .
<i>Office of Foreign Agricultural Relations</i>	LESLIE A. WHEELER, <i>Director</i> .
<i>Agricultural Adjustment Administration</i>	R. M. EVANS, <i>Administrator</i> .
<i>Bureau of Agricultural Chemistry and Engineering</i> .	HENRY G. KNIGHT, <i>Chief</i> .
<i>Bureau of Agricultural Economics</i>	H. R. TOLLEY, <i>Chief</i> .
<i>Agricultural Marketing Service</i>	C. W. KITCHEN, <i>Chief</i> .
<i>Bureau of Animal Industry</i>	JOHN R. MOHLER, <i>Chief</i> .
<i>Commodity Credit Corporation</i>	CARL B. ROBBINS, <i>President</i> .
<i>Commodity Exchange Administration</i>	J. W. T. DUVEL, <i>Chief</i> .
<i>Bureau of Dairy Industry</i>	O. E. REED, <i>Chief</i> .
<i>Bureau of Entomology and Plant Quarantine</i>	LEE A. STRONG, <i>Chief</i> .
<i>Farm Security Administration</i>	W. W. ALEXANDER, <i>Administrator</i> .
<i>Federal Crop Insurance Corporation</i>	LEROY K. SMITH, <i>Manager</i> .
<i>Federal Surplus Commodities Corporation</i>	MILO R. PERKINS, <i>President</i> .
<i>Food and Drug Administration</i>	WALTER G. CAMPBELL, <i>Chief</i> .
<i>Forest Service</i>	FERDINAND A. SILCOX, <i>Chief</i> .
<i>Bureau of Home Economics</i>	LOUISE STANLEY, <i>Chief</i> .
<i>Library</i>	CLARIBEL R. BARNETT, <i>Librarian</i> .
<i>Division of Marketing and Marketing Agreements</i> .	MILO R. PERKINS, <i>In Charge</i> .
<i>Bureau of Plant Industry</i>	E. C. AUCHTER, <i>Chief</i> .
<i>Rural Electrification Administration</i>	HARRY SLATTERY, <i>Administrator</i> .
<i>Soil Conservation Service</i>	H. H. BENNETT, <i>Chief</i> .
<i>Sugar Division</i>	JOSHUA BERNHARDT, <i>Chief</i> .
<i>Weather Bureau</i>	FRANCIS W. REICHELDERFER, <i>Chief</i> .